




International Journal of Advanced Research in Education and Technology (IJARETY)

Volume 13, Issue 2, March - April 2026

Impact Factor: 8.152



INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA



EcoTracker AI: An Intelligent Real-Time Environmental Monitoring and Analysis System

Bhagyashree Rathod

Department of Computer Science & Engineering, CMR Institute of Technology, Bengaluru, India

ABSTRACT: Environmental degradation, rising pollution levels, and climate change have made it increasingly critical to adopt smarter and more responsive environmental monitoring solutions. Conventional monitoring systems often rely on manual data collection, periodic inspections, and static reports, limiting the ability of individuals and organizations to respond quickly to changing environmental conditions. This paper presents EcoTracker AI, an intelligent environmental tracking system designed to deliver real-time monitoring and meaningful analysis through a single integrated digital platform. The system features a unified dashboard displaying live values of key environmental indicators—Air Quality Index (AQI), temperature, humidity, and noise levels—alongside a geo-tagged Live Map module for spatial awareness of pollution-affected zones. An AI-powered assistant enables natural language interaction with environmental data, removing the need for domain expertise. The full-stack architecture employs React.js on the frontend, Node.js and Express.js on the backend, and MongoDB for flexible data storage. Experimental results demonstrate that the system achieves average data-fetch response times below 200 ms and AI analysis execution times of approximately 250 ms under concurrent user loads. EcoTracker AI bridges the gap between raw environmental data and actionable human insight, supporting proactive decision-making and sustainable community awareness.

KEYWORDS: environmental monitoring, air quality index, real-time analytics, artificial intelligence, geospatial visualization, IoT, pollution detection, React.js, Node.js, MongoDB

I. INTRODUCTION

Environmental pollution and climate-related challenges represent some of the most pressing concerns of the twenty-first century. Rapid urbanization, industrial expansion, and increased vehicular emissions continue to degrade air, water, and acoustic environments worldwide [1]. Monitoring these changes in a timely, accurate, and accessible manner is essential for both regulatory compliance and individual health and safety decision-making.

Existing environmental monitoring solutions suffer from three principal limitations. First, data is typically gathered on fixed schedules—weekly, monthly, or quarterly—making it impossible to detect sudden pollution spikes or short-duration hazardous events. Second, outputs are presented as raw numerical tables or static PDF reports that are inaccessible to non-technical stakeholders. Third, systems are rarely integrated: air quality, weather, noise, and water quality data are siloed across different platforms, complicating holistic analysis [2].

The proliferation of low-cost sensors, open environmental APIs, and cloud computing infrastructure has created an opportunity to overcome these limitations. Researchers have proposed IoT-enabled sensor networks [3], machine learning-based anomaly detection [4], and GIS-driven spatial visualization [5] as complementary approaches. However, few systems combine all these capabilities into a single, user-accessible platform that also supports natural language interaction.

This paper presents EcoTracker AI, which addresses the above gap through the following contributions:

- A real-time, location-aware dashboard displaying AQI, temperature, humidity, and noise levels with automatic periodic refresh.
- A geo-tagged interactive map for spatial comparison of pollution hotspots across urban and peri-urban zones.
- An AI assistant supporting natural language queries about local environmental conditions and personalized eco-friendly recommendations.
- A predictive analytics module that forecasts short-term AQI trends based on historical sensor data.
- A role-based administrative panel for configuring alert thresholds, managing user accounts, and generating downloadable reports.
- A comprehensive feasibility analysis—technical, economic, operational, legal, and social—demonstrating viability for academic and smart-city deployment.

DOI:10.15680/IJARETY.2026.1302148

The remainder of this paper is structured as follows. Section II surveys related work. Section III describes the proposed system architecture and modules. Section IV details the software design, including UML diagrams and database schema. Section V presents implementation highlights. Section VI reports testing methodology and results. Section VII discusses performance metrics and comparative analysis. Section VIII concludes and outlines future directions.

II. RELATED WORK

A. Traditional Environmental Monitoring

Early environmental monitoring relied on manual field measurements and laboratory analysis, providing only periodic snapshots of conditions [2]. While regulatory compliance was achievable, the absence of continuous data made reactive management the norm, particularly in rapidly changing industrial zones. These limitations became critically apparent during pollution events of short duration that went undetected between scheduled sampling intervals.

B. IoT-Enabled Sensor Networks

The introduction of low-cost wireless sensor nodes enabled continuous, distributed data collection for parameters including particulate matter (PM_{2.5}), volatile organic compounds (VOCs), temperature, and humidity [3]. Studies demonstrated measurable improvements in spatial coverage and temporal resolution. However, interpreting the resulting data volumes remained a challenge for non-specialist users, as findings arrived as raw numerical streams without contextual explanation or actionable guidance.

C. Machine Learning for Pollution Analysis

Statistical and machine learning approaches—including ARIMA time-series models, random forests, and deep neural networks—have been applied to identify pollution trends, detect anomalies, and generate short-term forecasts [4], [7]. These methods outperform rule-based threshold systems in sensitivity and specificity, particularly under heterogeneous environmental conditions. Nonetheless, most published systems are research prototypes without a user-facing interface, limiting their practical adoption by non-expert stakeholders.

D. Geospatial Visualization

Geographic Information System (GIS) tools have proven effective for mapping pollution gradients and identifying hotspots [5]. Integration of live data feeds with interactive cartographic displays—using libraries such as Leaflet or the Google Maps API—allows users to observe spatial and temporal patterns simultaneously. Prior studies report that color-coded map overlays significantly reduce the time users need to identify high-risk zones compared with tabular data presentation, improving both response speed and comprehension accuracy.

E. Conversational Interfaces for Environmental Data

Research on chatbot-based environmental query systems indicates that natural language interfaces substantially improve user engagement and data accessibility for non-technical populations [6]. Conversational agents enable domain-agnostic queries about complex datasets, translating sensor readings into plain-language recommendations. However, combining conversational AI with real-time sensor data and geospatial visualization within a unified platform remains an open challenge—one that EcoTracker AI directly addresses by integrating all three capabilities into a single web application.

III. SYSTEM ARCHITECTURE

A. Overview

EcoTracker AI follows a three-tier client–server architecture comprising a React.js frontend, a Node.js/Express.js backend, and a MongoDB database layer, as depicted in Fig. 1. External environmental data APIs supply live AQI, weather, and pollution readings, while a dedicated analysis module processes incoming data streams for anomaly detection and trend forecasting. The architecture is designed for modularity, allowing independent scaling of each tier.




Fig. 1. High-level three-tier architecture of EcoTracker AI.

B. Frontend Layer

The frontend is implemented using React.js with a component-based architecture that enables smooth, single-page navigation across modules: Dashboard, Live Map, AI Insights, Pollution Analytics, Eco News, Eco Rewards, Reports, and AI Assistant. Recharts and Chart.js render time-series AQI graphs and seven-day weather trend charts. CSS color-coding (green / amber / red) provides immediate visual indication of safe, moderate, and hazardous environmental levels. The interface is fully responsive, supporting desktop, tablet, and mobile viewports through CSS Grid and Flexbox layouts.

C. Backend Layer

The backend server is built with Node.js (v16+) and Express.js, exposing RESTful API endpoints for user authentication, environmental data retrieval, AI query handling, alert generation, and report export. A Geolocation API automatically resolves the user's coordinates on login, routing subsequent data requests to location-specific external API endpoints. Data validation middleware rejects malformed or out-of-range sensor readings before persistence to the database, ensuring data integrity throughout the pipeline.

D. Data Analysis and AI Module

The analysis module performs three principal functions:

- Anomaly detection: compares incoming AQI, noise, and water quality readings against configurable safety thresholds; triggers alerts when limits are exceeded.
- Trend forecasting: applies time-series analysis to historical readings to estimate environmental conditions for the next seven days.
- Natural language response: the AI Assistant interprets free-text user queries, maps them to environmental data contexts, and returns plain-language explanations and eco-recommendations.

E. Database Layer

MongoDB (Community Edition / Atlas) stores user profiles, environmental data records, alerts, system logs, and eco-reward points in a document-oriented schema managed through Mongoose. The flexible document model accommodates the heterogeneous and variable-rate nature of environmental sensor data, supporting efficient querying for dashboard visualization and analytics. Indexes are applied on timestamp and location fields to optimize range queries for historical trend analysis.

F. Mapping Module

Interactive maps are rendered using the Leaflet library (or Google Maps API), displaying color-coded markers at geo-tagged measurement locations. Users can switch between "All Zones", "Hotspots", and "Safe Areas" map layers, enabling spatial comparison across regions in real time. Marker clusters are employed to handle dense sensor deployments without visual clutter, and pop-up tooltips display the most recent readings for each monitored location.

IV. SYSTEM DESIGN**A. Context Diagram**

The system context places EcoTracker AI as the central process interacting with four external entities: (1) the end user, who requests and receives environmental data and alerts; (2) the administrator, who monitors trends and configures system parameters; (3) external environmental data APIs, which supply real-time readings; and (4) the MongoDB database, which stores and retrieves all persistent records. Data flows are strictly validated at each boundary to prevent injection and unauthorized access.

B. Use Case Model

Two primary actors are identified:

General User: register/login; view real-time dashboard; check AQI and weather conditions; view 7-day trend charts; explore the live geo-tagged map; interact with the AI Assistant; view AI insights and pollution analysis; read Eco News; earn Eco Rewards; manage profile and notification preferences.

Administrator: secure admin login; monitor real-time and historical environmental data; view analytics and generate downloadable reports; manage user accounts; configure alert thresholds and data-refresh intervals; review system logs for anomalous activity.

C. Sequence Diagrams

Three key interaction flows are modeled as UML sequence diagrams:

- User viewing environmental data: User → Client App → Backend Server → External API → Backend (validate & store) → MongoDB → Client App (render dashboard with live metrics).
- Threshold alert generation: Backend fetches data → Analysis Module evaluates safety thresholds → if exceeded: generate alert → store in MongoDB → push notification to user interface.
- Admin login and log monitoring: Admin submits credentials → Backend verifies JWT against MongoDB → grant access to admin dashboard → retrieve historical logs and analytics → display trends and downloadable reports.

D. Database Schema

Table I summarizes the four primary MongoDB collections, their key fields, and field descriptions used to store and manage all persistent application data.

TABLE I PRIMARY DATABASE COLLECTIONS AND KEY FIELDS

Collection	Field	Description
3*User	user_id	Unique user identifier
	username	Display name
	location	Default geo-location
6*EnvData	data_id	Record identifier
	location	Measurement area
	AQI	Air Quality Index value
	pollution	Pollution level (float)
	weather	Weather condition string
	timestamp	UTC datetime of record
4*Alert	alert_id	Alert identifier
	data_id	Linked EnvData record
	severity	Low / Moderate / High
	time	Alert generation time
3*Admin	admin_id	Admin identifier
	username	Admin login name
	password	Bcrypt-hashed credential

V. IMPLEMENTATION**A. Technology Stack**

Table II summarizes the tools and technologies used across each layer of the EcoTracker AI system. The selection prioritizes open-source, actively maintained frameworks to minimize licensing costs and maximize community support.

TABLE II TECHNOLOGY STACK

Layer	Technology
Frontend	React.js, HTML5, CSS3, Recharts, Chart.js
Backend	Node.js (v16+), Express.js
Database	MongoDB, Mongoose ODM
Mapping	Leaflet / Google Maps API
Authentication	JWT (JSON Web Tokens)
Hosting	Railway / AWS
Version Control	Git

B. Real-Time Dashboard

The dashboard component polls the backend every 3,000 ms using a setInterval hook, randomly fluctuating AQI, temperature, and noise values within realistic bounds to simulate live sensor feeds during prototype evaluation. Color-coded status badges dynamically update based on computed AQI categories: Good (0–50), Moderate (51–100), Unhealthy (101–150), and Hazardous (>150). On production deployment, values are sourced directly from OpenWeatherMap and AQICN APIs keyed to the user's resolved geo-coordinates.

C. Pollution Analytics Module

The Pollution Analytics page incorporates a real-time noise meter using the Web Audio API (AudioContext, AnalyserNode), which captures microphone input, computes frequency-domain averages, and maps them to an approximate dB scale (30–120 dB). A simulated water quality panel displays pH (acceptable: 6.5–8.5), turbidity (NTU), and dissolved oxygen values for a representative urban lake, with color-coded status indicators showing safe, warning, and critical ranges.

D. AI Assistant

The chatbot frontend sends user queries to /api/chatbot via Axios. The backend constructs a structured prompt incorporating current AQI readings, location context, and historical trends, then forwards the enriched prompt to an external large-language model API. The response is returned as a plain-language environmental explanation. A local fallback function handles offline scenarios by keyword-matching queries to pre-defined responses covering AQI categories, weather patterns, water quality thresholds, waste reduction advice, and eco-facts.

E. Administrative Module

The admin panel enforces role-based access control (RBAC): only users with role: "admin" or role: "gov" stored in MongoDB are granted access after JWT verification. The panel provides tabular views of environmental trends, downloadable CSV/PDF reports via server-side generation, user management controls including account suspension and role assignment, and configurable alert threshold sliders for each monitored parameter. All administrative actions are logged with timestamps for audit purposes.

VI. TESTING**A. Functional Testing**

Each system module was tested individually (unit testing) and in combination (integration testing) through the browser interface. Key scenarios included: (1) correct rendering of dashboard metrics from external API data; (2) automatic data refresh on location change; (3) alert generation when AQI exceeded the configured threshold; (4) admin login with both

valid and invalid credentials; and (5) report generation for selected date ranges. All test cases were executed across Chrome, Firefox, and Safari to ensure cross-browser compatibility.

B. Non-Functional Testing

Performance testing was conducted with repeated concurrent API requests, confirming an average data-fetch response time below 200 ms. Reliability testing verified graceful degradation—cached data was displayed when external APIs were temporarily unavailable. Security testing confirmed that unauthorized users were blocked from admin endpoints by JWT middleware, invalid sensor inputs were rejected during data validation, and API keys were protected via server-side environment variables inaccessible to the client.

C. Test Case Summary

Table III presents the ten selected functional test cases, their descriptions, expected outcomes, and pass/fail results following systematic evaluation.

TABLE III SELECTED FUNCTIONAL TEST CASES

ID	Description	Expected Result	Status
TC01	Fetch environmental data for selected location	Dashboard updated	Pass
TC02	Invalid API response received	Data rejected	Pass
TC03	High pollution value triggers threshold	Alert displayed	Pass
TC04	Valid sensor data submitted	Stored in DB	Pass
TC05	Admin login with valid credentials	Dashboard shown	Pass
TC06	Admin login with invalid password	Error message	Pass
TC07	Report generation for date range	Report exported	Pass
TC08	Dashboard page reload	Latest data shown	Pass
TC09	Concurrent multi-user access	No conflict	Pass
TC10	Stored metrics rendered as charts	Graphs displayed	Pass

D. Boundary and Security Testing

Boundary cases tested included empty datasets, large historical datasets exceeding 10,000 records, rapid successive refresh requests at intervals below 500 ms, and simultaneous administrative configuration changes from multiple sessions. All cases passed without crashes or data corruption. Security tests additionally confirmed that direct database access was restricted to backend services only, preventing client-side query injection.

VII. RESULTS AND DISCUSSION

A. Performance Metrics

The system demonstrated strong runtime performance suitable for small-to-medium scale deployments. Table IV summarizes the key performance metrics observed during testing under concurrent user loads.

TABLE IV SYSTEM PERFORMANCE METRICS

Metric	Observed Value
Average API response time	< 200 ms
AI analysis execution time	≈ 250 ms per query
Dashboard render time (first load)	< 1.5 s
Concurrent user support	Stable (no degradation)

Data refresh interval	3,000 ms (configurable)
Alert generation latency	< 300 ms

B. Usability Observations

Color-coded AQI indicators (green/amber/red) and plain-language AI responses were found to be effective mechanisms for conveying environmental risk to non-technical users. The geo-tagged live map enabled rapid identification of pollution hotspots without requiring GIS expertise. The conversational AI assistant reduced the need for users to navigate multiple data screens, consolidating insights into a single interaction point. User testing feedback highlighted the Eco Rewards module as an effective behavioral incentive for sustained platform engagement and adoption of eco-friendly practices.

C. Comparison with Existing Approaches

Table V provides a structured comparison of EcoTracker AI against alternative environmental monitoring approaches across six key capability dimensions.

TABLE V COMPARISON WITH EXISTING APPROACHES

Feature	EcoTracker AI	Static Reports	Specialist GIS	Single-Param Sensors
Real-time updates	✓	✗	Partial	✓
Multi-parameter integration	✓	✓	✓	✗
No technical training needed	✓	✓	✗	✓
AI natural language interface	✓	✗	✗	✗
Geospatial hotspot mapping	✓	✗	✓	✗
Proactive threshold alerts	✓	✗	Partial	Partial

Unlike static reporting platforms, EcoTracker AI provides sub-second data updates and proactive threshold-based alerts. Unlike specialist GIS tools, it requires no technical training for operation. Unlike single-parameter sensors, it integrates AQI, weather, noise, and water quality into one coherent interface. The addition of an AI conversational layer distinguishes the system from prior integrated dashboards reported in the literature [3], [6].

VIII. CONCLUSION

This paper has presented EcoTracker AI, an intelligent, real-time environmental monitoring platform that integrates live data visualization, AI-powered analysis, geo-spatial mapping, threshold-based alerting, and natural language interaction within a unified web application. The system achieves sub-200 ms data-fetch response times and supports concurrent user access without performance degradation, making it suitable for deployment in academic institutions, smart-city initiatives, corporate sustainability programs, and community awareness campaigns.

All ten functional test cases passed across multiple browsers and device types. Boundary and security testing confirmed robustness under edge conditions and unauthorized access attempts. The open-source, modular technology stack facilitates cost-effective deployment and seamless future extension. The Eco Rewards mechanism introduces a novel behavioral dimension to environmental monitoring platforms, encouraging sustained user engagement and real-world adoption of sustainable practices.

EcoTracker AI demonstrates that bridging the gap between raw sensor data and actionable citizen insight is technically achievable within a low-cost, openly accessible web architecture. By unifying real-time monitoring, predictive analytics, conversational AI, and geospatial visualization, the platform represents a meaningful step toward intelligent, citizen-centered environmental governance.

IX. FUTURE WORK

Planned enhancements include: (1) integration of IoT hardware sensors for on-premise data collection independent of external APIs; (2) multi-language support targeting regional Indian languages (Hindi, Kannada, Telugu, Tamil) to broaden accessibility; (3) voice-based alert delivery for users with visual impairments or limited literacy; (4) satellite imagery analysis for large-scale land-use and air quality assessment; (5) cloud-native deployment on AWS or GCP with horizontal auto-scaling for high-traffic environments; and (6) expanded predictive analytics incorporating deep learning models for longer-horizon forecasting and cross-region pollution benchmarking.

X. ACKNOWLEDGMENT

The author thanks the faculty and peers of the Department of Computer Science & Engineering, CMR Institute Of Technology, for their guidance, constructive feedback, and encouragement throughout this research project.

REFERENCES

- [1] World Health Organization, "Air Quality Guidelines: Global Update," WHO Publications, Geneva, 2023.
- [2] European Environment Agency, "Environmental Indicators and Data Analysis," Publications Office of the European Union, Copenhagen, 2023
- [3] A. Kumar and S. Patel, "IoT-based real-time air quality monitoring system using low-cost sensors," IEEE Sensors Journal, vol. 21, no. 14, pp. 15823–15834, Jul. 2021.
- [4] L. Zhang, Y. Wang, and H. Chen, "Deep learning approaches for urban air quality prediction: A survey," Environmental Research, vol. 195, p. 110616, Apr. 2021.
- [5] M. R. Islam and T. Nakajima, "GIS-integrated environmental monitoring: Spatial analysis of urban pollution hotspots," Remote Sensing, vol. 13, no. 8, p. 1562, Apr. 2021.
- [6] S. Singh and R. Gupta, "Conversational AI for environmental awareness: Design and evaluation of a chatbot system," Proc. IEEE Int. Conf. on Computing, Communication and Security (ICCCS), pp. 1–6, 2022.
- [7] I. Goodfellow, Y. Bengio, and A. Courville, Deep Learning. Cambridge, MA: MIT Press, 2016.
- [8] National Aeronautics and Space Administration, "Earth Observation and Environmental Monitoring," NASA Technical Reports, 2023. [Online]. Available: <https://earthdata.nasa.gov>
- [9] OpenWeatherMap, "Weather and Environmental Data APIs Documentation," 2024. [Online]. Available : <https://openweathermap.org/api>
- [10] MongoDB, Inc., "MongoDB Database and Data Management," MongoDB Documentation, 2024. [Online]. Available: <https://www.mongodb.com/docs->

International Journal of Advanced Research in Education and Technology

ISSN: 2394-2975

Impact Factor: 8.152